

Examination of the Correlation of Distal End Parameters of the Humerus with Other Parameters

Humerus Distal Uç Parametrelerinin Diğer Parametrelerle İlişkisinin İncelenmesi

© Şeyma Toy, © Rukiye Sümeyye Bakıcı, © Necati Emre Şahin

Karabük University Faculty of Medicine, Department of Anatomy, Karabük, Turkey



Abstract

Objective: The aim of this study is to determine the correlation between all osteometric parameters obtained from the humerus and parameters obtained from the distal end of the humerus.

Materials and Methods: A total of 16 parameters were measured in 67 dry bones using the Image-J program.

Results: Groove for the ulnar nerve width (GUW) and the angle of the groove of the ulnar nerve (GUA) parameters were found to be higher on the left side and the difference was found to be statistically significant ($p<0.05$), GUA and intertubercular groove width were not significantly correlated with any parameter ($p>0.05$).

Conclusion: We believe that the results of the study will provide osteometric data of the humerus in distal end fracture treatment, comminuted fractures, and reconstructive surgery of the humerus, anthropological and forensic studies and increase the anatomical knowledge levels of experts interested in the region.

Keywords

Dry bone, distal end of the humerus, osteometry

Anahtar Kelimeler

Kuru kemik, humerus distal ucu, osteometri

Received/Geliş Tarihi : 27.01.2022

Accepted/Kabul Tarihi : 13.07.2022

doi:10.4274/meandros.galenos.2022.02439

Address for Correspondence/Yazışma Adresi:

Assoc. Prof. Şeyma Toy, PhD, Karabük
University Faculty of Medicine, Department of
Anatomy, Karabük, Turkey
Phone : +90 370 418 93 40
E-mail : seymatoy44@gmail.com
ORCID ID: orcid.org/0000-0002-6067-0087

Öz

Amaç: Bu çalışmanın amacı humerustan elde edilen tüm osteometrik parametreler ile humerus distal ucundan elde edilen parametreler arasındaki korelasyonu bulmaktır.

Gereç ve Yöntemler: Toplam 67 kuru kemikten, 16 parametre ölçümü Image-J programıyla yapıldı.

Bulgular: Sulcus ulnaris genişliği (GUW) ve sulcus ulnaris açısı (GUA) parametreleri sol tarafta daha yüksek bulundu ve aradaki fark istatistiksel olarak anlamlı bulundu ($p<0,05$), GUA ve sulcus intertubercularis genişliği herhangi bir parametre ile anlamlı olarak korelasyon göstermedi ($p>0,05$).

Sonuç: Çalışmanın sonuçlarının humerus distal uç kırıklarının tedavisi, parçalı kırıklar ve humerus rekonstrüktif cerrahisi, antropolojik ve adli çalışmalarda osteometrik veriler sağlayacağı ve bölgeyle ilgilenen uzmanların anatomik bilgi düzeylerini artıracacağı kanaatindeyiz.

Introduction

As in all biological structures, bone structure is driven by phylogenetic, structural, and functional factors (1,2).

The structure of the bone and its connection with its function is central to all inquiries that target or involve the bone. This includes deductive and inductive applications of skeletal biology both in medical science and also in archaeological or forensic sciences. Bone lengths and correlations have been studied by anthropologists, forensic scientists and orthopaedists for surgery and prosthetic design (3).

Distal humerus fractures account for 0.5-7% of all fractures, and approximately 30% of elbow fractures (4). Various implants are available for fracture patterns in the distal of humerus and these implants have been shaped according to the general anatomy knowledge of the elbow region (5).

The aim of this study is to investigate the relationship between humeral morphometry and humeral distal end parameters and other parameters of the humerus by digital measurements made on the dry bones of the humerus.

Materials and Methods

This study was initiated with 2021/797 numbered approval from Karabük University Ethics Committee (date: 24.12.2021). In the study, a total of 67 humerus bones, 35 left and 32 right, of unknown gender, were used from the bone collections in the anatomy laboratories of the medical faculty of the three universities. Deformed bones were not included in the study.

Morphometric Measurements

The bones placed on a measuring ruler were photographed with a professional digital camera (Nikon® - model D5300) parallel to the ground and from a height of 40 cm. The taken photos were transferred to Image-J (version 1.8 for Windows) program and they were calibrated. On the calibrated photos, 16 parameters were measured and recorded 2 times for each bone at different dates by the same observer.

Parameters;

The maximum length of the humerus (HML),
The vertical diameter of the humeral head (HVD),
Intertubercular groove width (GIW),
Deltoid tuberosity diameter (TDD),

Capitulum of the humerus width (CHW),
Trochlea of humerus width (THW),
Capitulum of humerus + trochlea of humerus transverse length,
Coronoid fossa width (TCW),
Radial fossa width,
The length between medial epicondyle and lateral epicondyle (FRW),
Groove for the ulnar nerve width (GUW),
The angle of the groove of the ulnar nerve (GUA),
Capitulum of humerus length (CHL),
Trochlea of humerus length (THL),
Olecranon fossa length (transverse plane) (FOL) and Olecranon fossa width (coronal plane) (FOW). All parameters showed in Figure 1.

Statistical Analyses

The data obtained in the study was analysed by using SPSS (Released: 2016, IBM SPSS Statistics for Windows, Version 24.0. Armonk, NY: IBM Corp.) software. In data evaluation, mean, standard deviation, minimum and maximum values were used as descriptive statistical methods. Normality of the continuous variables was evaluated with Shapiro-Wilk test. Two Sample t-test and Mann-Whitney




Figure 1. Parameters taken from the a) anterior, b) posterior, c) lateral, d) medial, e) and f) end of the humerus.

CHW: Capitulum of humerus width, EML: The length between medial epicondyle and lateral epicondyle, FCW: Coronoid fossa width, FOL: Olecranon fossa length (transverse plane), FRW: Radial fossa width, FOW: Olecranon fossa width (coronal plane), GIW: Intertubercular groove width, GUA: The angle of the groove of the ulnar nerve, GUW: Groove for the ulnar nerve width, HML: The maximum length of the humerus, HVD: The vertical diameter of the humeral head, TCW: Capitulum of humerus + trochlea of humerus transverse length, TDD: Deltoid tuberosity diameter, THL: Trochlea of humerus length, THW: Trochlea of humerus width

U test were used in the comparison of continuous variables between two groups. The linear correlation between the continuous variables was evaluated with Spearman and Pearson correlation test. P<0.05 level was considered as statistically significant.

In order to assess intra-observer precision and measurement accuracy, technical error of measurement (TEM), relative technical error of measurement (rTEM) and reliability coefficient (R) were calculated by using Microsoft Excel program.

RESULTS

The means of the first and second measurements were used in the study. TEM, rTEM, and R of the measurement parameters are shown in Table 1. TEM

Table 1. Intra-observer error analysis (TEM, rTEM, R) (n=67)

Parameters	TEM	rTEM (%)	R
GIW	0.11	12.49	0.83
FRW	0.07	5.56	0.88
THW	0.08	3.63	0.90
THL	0.10	3.82	0.91
FOW	0.06	3.53	0.93
TDD	0.06	3.10	0.94
FCW	0.06	3.47	0.94
G UW	0.07	5.18	0.94
CHW	0.05	2.82	0.95
FOL	0.06	2.30	0.96
CHL	0.05	2.18	0.97
TCW	0.04	0.95	0.98
GUA	1.69	1.44	0.98
HVD	0.05	1.05	0.98
HML	0.11	0.36	0.99
EML	0.05	0.84	0.99

TEM: Technical error of measurement, rTEM: Relative technical error of measurement, R: Reliability coefficient, GIW: Intertubercular groove width, FRW: Radial fossa width, THW: Trochlea of humerus width, THL: Trochlea of humerus length, FOW: Olecranon fossa width (coronal plane), TDD: Deltoid tuberosity diameter, FCW: Coronoid fossa width, G UW: Groove for the ulnar nerve width, CHW: Capitulum of humerus width, FOL: Olecranon fossa length (transverse plane), CHL: Capitulum of humerus length, TCW: Capitulum of humerus + trochlea of humerus transverse length, GUA: The angle of the groove of the ulnar nerve, HVD: The vertical diameter of the humeral head, HML: The maximum length of the humerus, EML: The length between medial epicondyle and lateral epicondyle

values are between 0.04 and 1.69. rTEM values are between 0.36% and 12.49%. R values are between 0.83 and 0.99, and the values are close to 1.

GIW, TDD, FCW, FOL, and FOW parameters were not distributed normally. No significant difference was found between the right and left side in these parameters (p>0.05) (Table 2). The other parameters were normally distributed; G UW and GUA parameters were found to be higher on the left side and the difference was found to be statistically significant (p<0.05) (Table 3).

Correlation Analysis Results

The correlation analysis is shown in Table 4. GUA and GIW were not significantly correlated with any parameter (p>0.05) (Table 4).

Table 2. Non-normally distributed parameters (n=67)

Parameters	Left	Right	p-value
GIW (cm)	0.89 (0.25-1.93)	0.92 (0.35-1.69)	0.252
TDD (cm)	2.17 (0.60-2.83)	2.20 (1.74-2.72)	0.497
FCW (cm)	1.84 (1.11-2.19)	1.73 (1.40-2.45)	0.250
FOL (cm)	2.76 (1.82-3.38)	2.85 (1.82-3.38)	0.748
FOW (cm)	1.91 (0.94-2.50)	1.93 (1.40-2.65)	0.713

GIW: Intertubercular groove width, TDD: Deltoid tuberosity diameter, FCW: Coronoid fossa width, FOL: Olecranon fossa length (transverse plane), FOW: Olecranon fossa width (coronal plane)

Table 3. Normally distributed parameters (n=67)

	Left	Right	p-value
FRW (cm)	1.401±0.213	1.384±0.243	0.765
THW (cm)	2.458±0.265	2.446±0.320	0.868
THL (cm)	2.724±0.335	2.740±0.394	0.859
G UW (cm)	1.559±0.301	1.328±0.292	0.003
CHW (cm)	1.962±0.251	1.886±0.233	0.207
CHL (cm)	2.404±0.267	2.411±0.351	0.932
TCW (cm)	4.478±0.417	4.403±0.386	0.459
GUA (°)	121.4 ±14.6	113.0±15.4	0.028
HVD (cm)	4.678±0.479	4.813±0.511	0.270
HML (cm)	32.37±2.68	31.99±2.29	0.534
EML (cm)	6.129±0.563	5.988±0.594	0.333

FRW: Radial fossa width, THW: Trochlea of humerus width, THL: Trochlea of humerus length, G UW: Groove for the ulnar nerve width, CHW: Capitulum of humerus width, CHL: Capitulum of humerus length, TCW: Capitulum of humerus + trochlea of humerus transverse length, GUA: The angle of the groove of the ulnar nerve, HVD: The vertical diameter of the humeral head, HML: The maximum length of the humerus, EML: The length between medial epicondyle and lateral epicondyle

Variables	HML	HVD	GIW	TDD	CHW	THW	TCW	FCW	FRW	EML	G UW	GUA	CHL	THL	FOL
1. HML (cm)															
2. HVD (cm)	0.675**														
3. GIW (cm)	0.189	0.157													
4. TDD (cm)	0.496**	0.598**	-0.286												
5. CHW (cm)	0.454**	0.419**	0.072	0.379**											
6. THW (cm)	0.526**	0.553**	0.084	0.421**	0.073										
7. TCW (cm)	0.716**	0.665**	0.168	0.503**	0.600**	0.730**									
8. FCW (cm)	0.399**	0.383**	0.193	0.299*	0.153	0.502**	0.423**								
9. FRW (cm)	0.252*	0.279*	0.086	0.342**	0.295*	0.159	0.277*	0.092							
10. EML (cm)	0.665**	0.577**	0.220	0.537**	0.444**	0.654**	0.798**	0.492**	0.158						
11. G UW (cm)	0.442**	0.372**	0.033	0.253	0.254*	0.325**	0.400**	0.341*	0.065	0.541**					
12. GUA (°)	-0.108	-0.218	-0.141	0.045	0.050	-0.199	-0.196	0.074	0.005	-0.240	0.084				
13. CHL (cm)	0.442**	0.539**	0.012	0.543**	0.553**	0.433**	0.665**	0.381**	0.241	0.610**	0.334**	-0.116			
14. THL (cm)	0.591**	0.631**	0.164	0.489**	0.406**	0.462**	0.591**	0.453**	0.069	0.608**	0.514**	-0.062	0.563**		
15. FOL (cm)	0.378**	0.345**	0.253	0.196	0.305*	0.278*	0.441**	0.185	0.278*	0.376**	0.148	-0.135	0.351**	0.234	
16. FOW (cm)	0.210	0.219	0.218	0.186	0.095	0.271	0.275*	0.277	0.098	0.301*	0.179	0.013	0.192	0.210	0.597**

**p<0.01, *p<0.05, HML: The maximum length of the humerus, HVD: The vertical diameter of the humeral head, GIW: Intertubercular groove width, TDD: Deltoid tuberosity diameter, CHW: Capitulum of humerus width, THW: Trochlea of humerus width, TCW: Capitulum of humerus + trochlea of humerus transverse length, FCW: Coronoid fossa width, FRW: Radial fossa width, EML: The length between medial epicondyle and lateral epicondyle, G UW: Groove for the ulnar nerve width, GUA: The angle of the groove of the ulnar nerve, CHL: Capitulum of humerus length, THL: Trochlea of humerus length, FOL: Olecranon fossa length (transverse plane), FOW: Olecranon fossa width (coronal plane)

Discussion

This study aimed to examine the correlation between measurement parameters of the distal end of humerus and all humerus parameters by making anthropometric measurements on dry bones of the humerus showed that HML parameter was associated with most of the distal end parameters. It was also found that parameters on distal end were generally correlated with each other. Another result was the finding that G UW and GUA parameters were higher on the left side when compared with the right side.

In this study, the measurements of which were repeated twice, TEM, rTEM, and R values of 16 parameters were calculated. R values of all parameters were found to be between 0.83 and 0.99, and they were interpreted to have almost perfect reliability (6,7). The highest rTEM percentage was calculated as 12.49% in GIW parameter. The reason why this parameter had high rTEM value is the fact that it had a mean lower than 1 cm (8).

Fractures to the long bones are common (9). In such cases, samples are needed to evaluate the methods used in implants. In the literature, various fixation experiments have been performed on these

specimens for clinical use based on screw-plate positioning. Knowing both the proximal and distal end of the humerus very well will increase the success in experiments such as biomechanical adjustment and durability on samples with various devices (10-13). Distal humeral prosthesis hemiarthroplasty used in comminuted distal humerus fractures should be designed to provide anatomical prerequisites and should be compatible with natural humeral anatomy. Having detailed knowledge of elbow anatomy will increase the design and success of new prostheses (14). At this point, the results obtained in this study will create an important anatomical resource.

Segmental studies of the humerus, especially in forensic medicine and archeology, there are studies that estimate the height from the humerus by using various formulas to determine the height of the people in the population (15,16). We think that HML will be a reference for such studies.

Kastamoni et al. (17) stated that G UW was positively significantly correlated with humeral retroversion angle and HML and negatively significantly correlated with olecranon fossa depth. Since the parameters of humeral retroversion angle and olecranon fossa

depth were not included in our study, no inferences were made in our study on this. Although Vettivel et al. (18) found positively significant correlation with GUA and HML in their study, no significant correlation was found in the present study. Wafae et al. (19) stated that intertubercular groove morphology was not correlated with humeral morphometry. Measurements were made from the middle level of intertubercular groove both in this study and in Wafae et al.'s (19) study, while measurement was made from the proximal of the groove in Kastamoni et al.'s (17) study. We think that the difference in the results may be due to this reason.

Kastamoni et al. (17) concluded that there were no significant differences between olecranon fossa width and HML, EML, and THW parameters. On the other hand, a statistically significant correlation was found between FOW and EML in our study.

There are studies in literature which report correlation between HML and EML (20-22). There are also studies reporting a significant correlation between HML and THW parameters (22, 23). Similar results found in our study support the existing literature. While Kastamoni et al. (17) stated that HML parameter is not significantly correlated with EML and THW parameters, this result is not consistent with the results of our study and other studies. On the other hand, Kastamoni et al. (17) found statistically significant positive correlation between EML and THW and this result was found to be consistent with the results of our study.

There are studies in literature which show statistically significant positive correlation between HML and HVD parameters and this result is consistent with the results of our study (21,22). Similar to the results of our study, Çini and Arı (21) found significant positive correlation between HML and CHW parameters. However, although they found significant correlation between HML and GUA parameters, no significant correlation was found between these parameters in our study.

de Queiroz Chaves et al. (24) conducted by using the computed tomography images of the right sides of 30 individuals, they found mean GUA parameter as 92.7 ± 10 . In our study, this parameter was found as $113.0 \pm 15.4^\circ$ on the right side. The fact that we had no recorded information about general health state since

our study was conducted on dry bones, uncertain gender distribution and age, demographic differences and differences in sample size may be the reasons for the difference between studies. GUA parameter was found to be higher on the left side when compared with the right side in our study and the difference was found to be statistically significant. Since the same person was not compared bilaterally in our study, this difference was weak in terms of evidence. This parameter was not found to be correlated with any of our parameters. It has been stated in literature that bone structure is important in cubital tunnel syndrome factors (25). GUA is also the parameter that can determine the shape of cubital tunnel and we believe that it is clinically important.

This study has some limitations. Not having information about the age and gender of the dry bones used, about general health state and about whether the right and left bones belonged to same people are important limitations of the study.

Conclusion

Nevertheless, the anatomical information found in this highly reliable study will guide clinicians in studies evaluating humeral segments, especially in the treatment of elbow instability, the treatment and reconstruction of distal humeral fractures. It will help the multidisciplinary team to produce a more suitable prosthesis both biomechanically and kinesiology by correlating it with radiological data. The results of this study will contribute to studies in the fields of anatomy, biomechanics, forensic sciences, and anthropology.

Acknowledgment

We are grateful to Düzce University Department of Anatomy and Bolu Abant İzzet Baysal University Department of Anatomy for allowing us to work on the bone collections.

Ethics

Ethics Committee Approval: This study was initiated with 2021/797 numbered approval from Karabük University Ethics Committee (date: 24.12.2021).

Informed Consent: Informed consent is not required.

Peer-review: Externally and internally peer-reviewed.

Authorship Contributions

Concept: R.S.B., Design: R.S.B., Data Collection or Processing: Ş.T., R.S.B., N.E.Ş., Analysis or Interpretation: Ş.T., R.S.B., N.E.Ş., Literature Search: Ş.T., R.S.B., N.E.Ş., Writing: Ş.T., R.S.B., N.E.Ş.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

- Gould SJ. The structure of evolutionary theory. Harvard University Press; 2002.
- Cubo J. Pattern and process in constructional morphology. *Evol Dev* 2004; 6: 131-3.
- Pietak A, Ma S, Beck CW, Stringer MD. Fundamental ratios and logarithmic periodicity in human limb bones. *J Anat* 2013; 222: 526-37.
- Galano GJ, Ahmad CS, Levine WN. Current treatment strategies for bicolumnar distal humerus fractures. *J Am Acad Orthop Surg* 2010; 18: 20-30.
- Narayana AS, Aradhyula H. The Morphometric Study Of Distal End Of Humerus. *Paripex - Indian Journal Of Research*. 2018; 7: 136-7.
- Jamayah H, Geeta A, Safiza M, Khor G, Wong N, Kee C, et al. Reliability, technical error of measurements and validity of length and weight measurements for children under two years old in Malaysia. *Med J Malaysia*. 2010; 65:131-7.
- Perini TA, de Oliveira GL, Ornellas JdS, de Oliveira FP. Technical error of measurement in anthropometry. *Rev Bras Med Esporte* 2005; 11: 86-90.
- Bakici RS, Oner Z, Oner S. The analysis of sacrum and coccyx length measured with computerized tomography images depending on sex. *Egypt J Forensic Sci* 2021; 11: 14.
- Öztürk K, Dursun A, Ayazoğlu M, Kastamoni Y. An Investigation of the Nutrient Foramen in the Long Bones of the Upper and Lower Limbs in Turkish Population. *Med Records* 2022; 4:179-86.
- Scolaro JA, Hsu JE, Svach DJ, Mehta S. Plate selection for fixation of extra-articular distal humerus fractures: A biomechanical comparison of three different implants. *Injury* 2014; 45: 2040-4.
- Wilson DJ, Scully WF, Min KS, Harmon TA, Eichinger JK, Arrington ED. Biomechanical analysis of intramedullary vs. superior plate fixation of transverse midshaft clavicle fractures. *J Shoulder Elbow Surg* 2016; 25: 949-53.
- Hackl M, Wegmann K, Taibah S, Burkhart KJ, Scaal M, Müller LP. Peri-implant failure in dual plating of the distal humerus. A biomechanical analysis with regard to screw and plate positioning. *Injury* 2015; 46: 2142-5.
- Mighell MA, Stephens B, Stone GP, Cottrell BJ. Distal Humerus Fractures: Open Reduction Internal Fixation. *Hand Clin* 2015; 31: 591-604.
- Heijink A, Wagener ML, de Vos MJ, Eygendaal D. Distal humerus prosthetic hemiarthroplasty: midterm results. *Strategies Trauma Limb Reconstr* 2015; 1: 101-8.
- Vinaykumar K, Asha K, Bindurani M, Kavyashree A, Suresh N. Estimation of Humeral Length from its Proximal and Distal Fragments in South Indian Population. *Medico-Legal Update*. 2015; 15: 55.
- Kantha L, Kulkarni R. Estimation of Total Length Of Humerus From Its Fragments In South Indian Population. *Int J Anat Res* 2014; 2: 213-20.
- Kastamoni Y, Yazan H, Dursun A, Öztürk K, Özgel Ö, Albay S. Morphometric Evaluation of Humerus and Its Clinical Significance. *Süleyman Demirel University Journal of Health Science*. 2021; 12: 77-85.
- Vettivel S, Indrasingh I, Chandi G, Chandi S. Variations in the intertubercular sulcus of the humerus related to handedness. *J Anat* 1992; 180: 321-6.
- Wafae N, Santamaría LEA, Vitor L, Pereira LA, Ruiz CR, Wafae GC. Morphometry of the human bicipital groove (sulcus intertubercularis). *J Shoulder Elbow Surg* 2010; 19: 65-8.
- Hamzehtofigh M, Bayat P, Rahimifar R. Sex Determination from the Humerus Bone in Iranian Cases. *Int J Morphol* 2019; 37: 1370-4.
- Çini NT, Arı İ. Estimation of total bone length of the late Byzantine humerus. *Archaeol Anthropol Sci* 2021; 13: 1-8.
- Bhusaraddi PS, Shinde V, Khona P. Right humerus; An equation to estimate the length from its fragments. *IP Indian J Anat Surg Head Neck Brain* 2019; 5: 101-4.
- Giannicola G, Scacchi M, Sedati P, Gumina S. Anatomical variations of the trochlear notch angle: MRI analysis of 78 elbows. *Musculoskelet Surg* 2016; 100: 89-95.
- de Queiroz Chaves L, de Souza Fonseca GV, da Silva FHP, Acioly MA. Osseous morphology of the medial epicondyle: an anatomoradiological study with potential clinical implications. *Surg Radiol Anat* 2021; 43: 713-20.
- Lee SK, Hwang SY, Kim SG, Choy WS. Analysis of the Anatomical Factors Associated with Cubital Tunnel Syndrome. *Orthop Traumatol Surg Res* 2020; 106: 743-9.